

**Suspected Upper Extremity Deep Vein Thrombosis
EVIDENCE TABLE**

Reference	Study Type	Patients/ Events	Study Objective (Purpose of Study)	Study Results	Strength of Evidence
1. Joffe HV, Goldhaber SZ. Upper-extremity deep vein thrombosis. <i>Circulation</i> 2002; 106(14):1874-1880.	12	N/A	To review risk factors, diagnostic options, treatment alternatives, and prophylaxis regimens of upper extremity deep vein thrombosis (UEDVT).	<ul style="list-style-type: none"> Although aggressive multimodal treatment is generally recommended for patients with primary UEDVT, this practice should be evaluated critically with prospective clinical trials. Preliminary studies suggest that US (without pharmacotherapy) may accelerate thrombolysis by enhancing enzymatic fibrinolysis and mechanically disrupting the thrombus. Significantly lower doses of thrombolytics may be effective when used in combination with US, thereby reducing bleeding complications. Further research is needed to evaluate the safety and efficacy of this novel treatment approach. 	4
2. Weissleder H, Weissleder R. Lymphedema: evaluation of qualitative and quantitative lymphoscintigraphy in 238 patients. <i>Radiology</i> 1988; 167(3):729-735.	10	238	To evaluate qualitative and quantitative lymphoscintigraphy in 128 patients with primary lymphedema, in 91 patients with secondary lymphedema, and in 19 healthy volunteers.	With qualitative interpretation alone, the diagnosis of lymphedema was established in 216 of 308 extremities (70.1%). Quantitative parameters derived from clearance data showed abnormal lymphatic function in all 308 extremities. Whereas qualitative lymphoscintigraphy allows the characterization of lymphatic morphology, quantitative lymphoscintigraphy is very accurate in detection of incipient lymphedema.	2
3. Abdullah BJ, Mohammad N, Sangkar JV, et al. Incidence of upper limb venous thrombosis associated with peripherally inserted central catheters (PICC). <i>Br J Radiol</i> 2005; 78(931):596-600.	3a	26	To prospectively determine the incidence of venous thrombosis in the upper limbs in patients with peripherally inserted central catheters (PICC).	<ul style="list-style-type: none"> No statistical correlation between the site of insertion of the PICC and the location of venous thrombosis. No observed correlation between the occurrence of venous thrombosis with the patient's history of hypertension, hypercholesterolaemia, coronary artery disease, diabetes mellitus, cardiac insufficiency, smoking or cancer. No statistical correlation with the size of the catheter. PICC are associated with a significant risk of UEDVT. 	3

**Suspected Upper Extremity Deep Vein Thrombosis
EVIDENCE TABLE**

Reference	Study Type	Patients/ Events	Study Objective (Purpose of Study)	Study Results	Strength of Evidence
4. Baarslag HJ, Koopman MM, Reekers JA, van Beek EJ. Diagnosis and management of deep vein thrombosis of the upper extremity: a review. <i>Eur Radiol</i> 2004; 14(7):1263-1274.	12	N/A	To review the clinical background, imaging modalities that may be employed, treatment options and outcome of patients with UEDVT.	No results reported.	4
5. Prandoni P, Polistena P, Bernardi E, et al. Upper-extremity deep vein thrombosis. Risk factors, diagnosis, and complications. <i>Arch Intern Med</i> 1997; 157(1):57-62.	9	58	To identify the clinical and laboratory parameters associated with UEDVT, to assess the diagnostic accuracy of ultrasonographic methods for its detection, and to establish the frequency of both early and late complications.	<ul style="list-style-type: none"> • Central venous catheters, thrombophilic states, and a previous leg vein thrombosis were statistically significantly associated with UEDVT. • Sensitivity and specificity of compression ultrasonography (96% and 93.5%, respectively) and color flow Doppler imaging (100% and 93%, respectively) were comparable and better than those of Doppler ultrasonography (81% and 77%, respectively). • Objective findings suggestive of a pulmonary embolism were recorded in 36% of the patients with UEDVT. After a mean follow-up of 2 years, two patients with UEDVT experienced recurrent thromboembolic events, and four had post-thrombotic sequelae. 	2
6. Schmittling ZC, McLafferty RB, Bohannon WT, Ramsey DE, Hodgson KJ. Characterization and probability of upper extremity deep venous thrombosis. <i>Ann Vasc Surg</i> 2004; 18(5):552-557.	13	177 upper extremities of arms	Retrospective review of patients who underwent color-flow duplex scanning for clinically suspected acute UEDVT. To characterize patient demographics, risk factors, and anatomic distribution of UEDVT to develop a probability model for diagnosis.	<ul style="list-style-type: none"> • Color-flow duplex scanning identified acute UEDVT in 53 (30%) of the arms examined with deep system involvement in 40 (23%). Of the upper extremities affected, the subclavian was involved in 64%, the axillary in 25%, the internal jugular in 32%, the brachial in 36%, the cephalic in 32%, and the basilic in 47%. • Use of this model can help focus clinical suspicion, improve color-flow duplex utilization, and provide timely treatment with anticoagulation. 	2
7. Spencer FA, Emery C, Lessard D, Goldberg RJ. Upper extremity deep vein thrombosis: a community-based perspective. <i>Am J Med</i> 2007; 120(8):678-684.	13	483	Review medical records to examine the magnitude, risk factors, management strategies, and outcomes in a population-based investigation of patients with upper, as compared with lower, extremity DVT diagnosed in 1999.	Patients with UEDVT represent a clinically important patient population in the community setting. Risk factors, occurrence of pulmonary embolism, and timing and location of venous thromboembolism recurrence differ between patients with upper as compared with lower extremity DVT.	2

**Suspected Upper Extremity Deep Vein Thrombosis
EVIDENCE TABLE**

Reference	Study Type	Patients/ Events	Study Objective (Purpose of Study)	Study Results	Strength of Evidence
8. Baarslag HJ, van Beek EJ, Koopman MM, Reekers JA. Prospective study of color duplex ultrasonography compared with contrast venography in patients suspected of having deep venous thrombosis of the upper extremities. <i>Ann Intern Med</i> 2002; 136(12):865-872.	9	126	Prospective study of duplex US compared with venography to determine the accuracy of duplex US for diagnosis of UEDVT.	<ul style="list-style-type: none"> Sensitivity and specificity of duplex US and venography were 82% (95% CI, 70% to 93%) and 82% (CI, 72% to 92%), respectively. Duplex US may be the method of choice for initial diagnosis of patients with suspected UEDVT. However, in patients with isolated flow abnormalities, contrast venography should be performed. 	2
9. Shebel ND, Marin A. Effort thrombosis (Paget-Schroetter syndrome) in active young adults: current concepts in diagnosis and treatment. <i>J Vasc Nurs</i> 2006; 24(4):116-126.	12	N/A	To review the prevalence, differential diagnosis, diagnostic modalities and medical and surgical interventions used to treat Paget-Schroetter.	Patients presenting with possible symptoms of Paget-Schroetter syndrome should be screened with an objective diagnostic test. Venous duplex imaging can identify acute thrombosis, but venography is necessary to confirm the diagnosis of Paget-Schroetter syndrome. Catheter-directed thrombolysis should be initiated in acute DVT to reduce the clot burden, and it offers a better long-term outcome.	3
10. Knudson GJ, Wiedmeyer DA, Erickson SJ, et al. Color Doppler sonographic imaging in the assessment of upper-extremity deep venous thrombosis. <i>AJR</i> 1990; 154(2):399-403.	10	91 patients 130 extremities	To determine the accuracy of color Doppler flow imaging in the assessment of UEDVT.	Sensitivity of the Doppler studies was 78% and the specificity was 92%. Color Doppler flow imaging is an accurate, noninvasive method for the evaluation of UEDVT. Other imaging tests may be required when the color Doppler study is negative and central venous thrombosis is suspected.	2
11. Mustafa S, Stein PD, Patel KC, Otten TR, Holmes R, Silbergleit A. Upper extremity deep venous thrombosis. <i>Chest</i> 2003; 123(6):1953-1956.	13	44,136	Retrospective review to determine the prevalence of symptomatic UEDVT and its association with symptomatic acute pulmonary embolism (PE) in a community teaching hospital.	Symptomatic UEDVT is not uncommon in hospitalized patients. Symptomatic PE resulting from UEDVT was not observed in these patients, all of whom were treated with anticoagulants.	2
12. Patel MC, Berman LH, Moss HA, McPherson SJ. Subclavian and internal jugular veins at Doppler US: abnormal cardiac pulsatility and respiratory phasicity as a predictor of complete central occlusion. <i>Radiology</i> 1999; 211(2):579-583.	9	21	To analyze changes in venous Doppler waveforms of damped or diminished cardiac pulsatility and respiratory phasicity. US findings were compared with phlebographic findings.	US can be used to establish the presence or absence of thrombosis in the distal portion of the brachiocephalic or subclavian veins, which are inaccessible to direct insonation.	3

**Suspected Upper Extremity Deep Vein Thrombosis
EVIDENCE TABLE**

Reference	Study Type	Patients/ Events	Study Objective (Purpose of Study)	Study Results	Strength of Evidence
13. Ong B, Gibbs H, Catchpole I, Hetherington R, Harper J. Peripherally inserted central catheters and upper extremity deep vein thrombosis. <i>Australas Radiol</i> 2006; 50(5):451-454.	13	317 scans	Retrospective study of all UEDVT duplex scans was carried out to determine the incidence and risk factors for venous thrombosis in patients with a PICC.	<ul style="list-style-type: none"> • Three main risk factors identified were; presence of a central line, malignancy and administration of chemotherapy. PICC were the most common central line present. • Symptomatic thrombosis occurred in 7% of PICC inserted for chemotherapy compared with 1% of PICC inserted for other reasons. 	3
14. Agarwal AK, Patel BM, Haddad NJ. Central vein stenosis: a nephrologist's perspective. <i>Semin Dial</i> 2007; 20(1):53-62.	12	N/A	To review management of central vein stenosis.	Endovascular interventions are the mainstay of management of central vein stenosis. Percutaneous angioplasty and stent placement for elastic and recurring lesions can restore the functionality of the vascular access, at least temporarily.	4
15. Lam EY, Giswold ME, Moneta GL. Venous and Lymphatic Disease. In: Brunicaardi FC, Andersen DK, Billiar TR, et al., eds. <i>Schwartz's Principles of Surgery</i> . 8th ed: McGraw-Hill; 2005.	15	N/A	Textbook.	N/A	N/A
16. Hingorani AP, Ascher E, Markevich N, et al. Prospective evaluation of combined upper and lower extremity DVT. <i>Vasc Endovascular Surg</i> 2006; 40(2):131-134.	13	227	Prospective study to evaluate combined upper and lower extremity DVT.	211 (93%) of 227 patients had lower extremity studies; 45 of these 211 (21%) had acute lower extremity DVT by duplex examination in addition to the UEDVT. 22 of 211 patients had bilateral lower extremity DVT, and 8 patients were found to have chronic thrombosis of lower extremity veins. Of the patients with bilateral UEDVT, there were 3 with bilateral lower extremity acute DVT. Finally, 8 of the remaining 166 patients (5%) with originally negative lower extremity studies were found to develop a thrombosis at a later date.	2
17. Do B, Mari C, Biswal S, Kalinyak J, Quon A, Gambhir SS. Diagnosis of aseptic deep venous thrombosis of the upper extremity in a cancer patient using fluorine-18 fluorodeoxyglucose positron emission tomography/computerized tomography (FDG PET/CT). <i>Ann Nucl Med</i> 2006; 20(2):151-155.	14	1	To describe a patient with a history of recurrent squamous cell carcinoma of the tongue and abnormal FDG uptake in the left arm during a re-staging FDG-PET/CT.	After revision of the patient's clinical history, tests and physical exam, the abnormal FDG uptake was found to correspond to an extensive aseptic UEDVT.	4

Suspected Upper Extremity Deep Vein Thrombosis
EVIDENCE TABLE

Reference	Study Type	Patients/ Events	Study Objective (Purpose of Study)	Study Results	Strength of Evidence
18. Gloviczki P, Calcagno D, Schirger A, et al. Noninvasive evaluation of the swollen extremity: experiences with 190 lymphoscintigraphic examinations. <i>J Vasc Surg</i> 1989; 9(5):683-689; discussion 690.	10	115 patients 190 extremities	Lymphoscintigraphy performed with Tc-99m labeled antimony trisulfide colloid was used as a noninvasive diagnostic examination to evaluate the lymphatic circulation in extremities.	<ul style="list-style-type: none"> Semiquantitative evaluation of the lymphatic drainage and visual interpretation of the image patterns were reliable to differentiate lymphedema from edemas of other origin (sensitivity: 92%, specificity: 100%). Lymphoscintigraphy is safe and reliable and has no side effects. It should replace contrast lymphangiography in the routine evaluation of the swollen extremity. 	2
19. Wang YF, Cherng SC, Chiu JS, Su YC, Sheu YT. Application of upper extremity radionuclide venography as a diagnostic approach for Port-A catheter thrombosis. <i>J Chin Med Assoc</i> 2006; 69(8):358-363.	10	14	To investigate the role of upper extremity radionuclide venography as a potential diagnostic modality in the assessment of venous thrombosis associated with a Port-A catheter.	Upper extremity radionuclide venography is an easily performed and effective method for diagnosing Port-A catheter thrombosis in clinical practice.	3
20. Baxter GM, Kincaid W, Jeffrey RF, Millar GM, Porteous C, Morley P. Comparison of colour Doppler ultrasound with venography in the diagnosis of axillary and subclavian vein thrombosis. <i>Br J Radiol</i> 1991; 64(765):777-781.	9	66	Prospective study to compare bilateral venography with US for detection of DVT in the upper venous system in children with acute lymphoblastic leukemia.	<ul style="list-style-type: none"> DVT occurred in 29% (19/66) patients. While 15/19 DVT were detected by venography (sensitivity 79%), only 7/19 were detected by US (sensitivity 37%). US is insensitive for DVT in the central upper venous system but may be more sensitive than venography in the jugular veins. A combination of both venography and US is required for screening for DVT in the upper venous system. 	1
21. Koksoy C, Kuzu A, Kutlay J, Erden I, Ozcan H, Ergin K. The diagnostic value of colour Doppler ultrasound in central venous catheter related thrombosis. <i>Clin Radiol</i> 1995; 50(10):687-689.	9	44	Prospective study to establish the correlation between color Doppler US observations and venography in detecting catheter-related thrombosis. Criteria used to show the presence of catheter-related thrombosis included visualization of thrombus (T), absence of spontaneous flow (S), absence of phasicity of flow with respiration (P), incompressibility of the vein with probe pressure (C) and visualization of increased venous collaterals (V).	Analyzing each variable separately, P had the highest sensitivity (94%) and S had the highest specificity (100%). The best combinations for diagnosis were T+P, P+S, and T+P+S with a sensitivity of 94% and specificity of 88%. Overall diagnostic value of color Doppler US had a sensitivity of 94% and specificity of 96%.	2
22. Weissleder R, Elizondo G, Stark DD. Sonographic diagnosis of subclavian and internal jugular vein thrombosis. <i>J Ultrasound Med</i> 1987; 6(10):577-587.	14	7	Patients with internal jugular and/or subclavian vein thrombosis were studied with real-time sonography and venography. High-resolution real-time sonography was used to tabulate morphologic parameters and physiologic parameters.	Two new signs of thrombosis, the absence of the beating venous valve and the "cut-off sign," are particularly useful in the sonographic diagnosis of internal jugular vein thrombosis.	4

**Suspected Upper Extremity Deep Vein Thrombosis
EVIDENCE TABLE**

Reference	Study Type	Patients/ Events	Study Objective (Purpose of Study)	Study Results	Strength of Evidence
23. Chin EE, Zimmerman PT, Grant EG. Sonographic evaluation of upper extremity deep venous thrombosis. <i>J Ultrasound Med</i> 2005; 24(6):829-838; quiz 839-840.	13	N/A	To review the techniques of performing an upper extremity Doppler examination, in addition to illustrating the sonographic appearances of acute and chronic UEDVT.	Color Doppler sonography is accurate in the diagnosis of UEDVT. However, in cases of equivocal Doppler findings, or when the sonographic findings are normal but clinical suspicion for central venous thrombosis is high, MRI or contrast venography is necessary for further evaluation.	3
24. Haire WD, Lynch TG, Lund GB, Lieberman RP, Edney JA. Limitations of magnetic resonance imaging and ultrasound-directed (duplex) scanning in the diagnosis of subclavian vein thrombosis. <i>J Vasc Surg</i> 1991; 13(3):391-397.	9	43 phlebograms	Correlate the results of phlebograms with duplex and MRI scans of the thoracic veins to determine the potential role of MRI and duplex scanning in the diagnosis of catheter-induced subclavian vein thrombosis.	<ul style="list-style-type: none"> • Duplex scans detected 6 of 11 occlusions, whereas MRI detected 4 of the 5 occlusions scanned. • Of 14 nonocclusive thrombi seen on phlebography, duplex scans correctly identified 8. • MRI was done on 8 nonocclusive thrombi but identified only 2. • Short occlusions of the proximal portion of the left subclavian vein were often undetected by duplex scanning but occasionally seen with MRI. • MRI is highly reliable in ruling out the presence of a thrombotic process in the subclavian vein, but it may on occasion fail to detect the presence of subclavian thrombi. 	3
25. Svensson WE, Mortimer PS, Tohno E, Cosgrove DO. Colour Doppler demonstrates venous flow abnormalities in breast cancer patients with chronic arm swelling. <i>Eur J Cancer</i> 1994; 30A(5):657-660.	10	81	Patients with chronic arm swelling related to their breast cancer treatment were examined with color Doppler imaging. Findings are reported.	Over half (57%) of the patients had evidence of venous outflow obstruction and a further 14% had signs of venous “congestion”. Only 30% of the swollen arms had normal venous outflow. The venous systems of the contralateral non-swollen arms were all normal as were both arms in 28 control patients who had similar treatment but had not developed arm swelling. Findings suggest that venous outflow obstruction is an important contributory factor in the pathophysiology of arm swelling following breast cancer treatment.	2
26. Weber TM, Lockhart ME, Robbin ML. Upper extremity venous Doppler ultrasound. <i>Radiol Clin North Am</i> 2007; 45(3):513-524.	12	N/A	To review the role of US in the evaluation of UEDVT.	US provides an accurate, rapid, low-cost, portable, noninvasive method for screening, mapping, and surveillance of the UEDVT system.	4

**Suspected Upper Extremity Deep Vein Thrombosis
EVIDENCE TABLE**

Reference	Study Type	Patients/ Events	Study Objective (Purpose of Study)	Study Results	Strength of Evidence
27. Grassi CJ, Polak JF. Axillary and subclavian venous thrombosis: follow-up evaluation with color Doppler flow US and venography. <i>Radiology</i> 1990; 175(3):651-654.	9	13	Real-time US and color Doppler flow mapping were used to determine prospectively whether UEDVT could be imaged as accurately as with conventional contrast venography.	Color Doppler flow US is useful in evaluating UEDVT and might be preferable to venography for follow-up because the discomfort of injection and risk of contrast material-induced phlebitis are eliminated.	3
28. Gaitini D, Beck-Razi N, Haim N, Brenner B. Prevalence of upper extremity deep venous thrombosis diagnosed by color Doppler duplex sonography in cancer patients with central venous catheters. <i>J Ultrasound Med</i> 2006; 25(10):1297-1303.	12	N/A	To review the literature concerning UEDVT diagnosed by color Doppler duplex sonography in cancer patients with indwelling central venous catheters.	Color Doppler duplex sonography is an accurate examination for the diagnosis of UEDVT, with sensitivity ranging from 78% to 100% and specificity ranging from 82% to 100%.	3
29. Haire WD, Lynch TG, Lieberman RP, Lund GB, Edney JA. Utility of duplex ultrasound in the diagnosis of asymptomatic catheter-induced subclavian vein thrombosis. <i>J Ultrasound Med</i> 1991; 10(9):493-496.	10	32 subclavian catheters	Prospective study to determine the usefulness of duplex US in the diagnosis of asymptomatic catheter-induced subclavian vein thrombosis.	16 arm venograms were normal and all gave normal duplex scans. 11 venograms demonstrated nonocclusive mural thrombi. Only 3 of these were seen with duplex US. Five totally occlusive thrombi were seen on venography, of which only 2 were detected with duplex sonography. The insensitivity of this technique to asymptomatic subclavian thrombi limits its usefulness as a screening tool.	3
30. Ho VB, Corse WR, Hood MD, Rowedder AM. Magnetic resonance angiography of the thoracic vessels. <i>Magnetic Resonance Imaging Clinics of North America</i> 2004; 12(4):727-747.	12	N/A	To review MR techniques used in the evaluation of the thoracic vessels.	Most thoracic vascular lesions can be well evaluated if the appropriate combination of pulse sequences is performed.	4
31. Hansen ME, Spritzer CE, Sostman HD. Assessing the patency of mediastinal and thoracic inlet veins: value of MR imaging. <i>AJR</i> 1990; 155(6):1177-1182.	13	31	Review hospital records and MR studies to assess the accuracy of MRI in the evaluation of patients with suspected thoracic venous obstruction.	MR correctly identified abnormality in 32 of 34 vessels (sensitivity 94%). In all cases, a negative MR study was found to be a true negative (no false positives in 42 normal vessels; specificity, 100%).	2
32. Baarslag HJ, Van Beek EJ, Reekers JA. Magnetic resonance venography in consecutive patients with suspected deep vein thrombosis of the upper extremity: initial experience. <i>Acta Radiol</i> 2004; 45(1):38-43.	9	44	To assess the feasibility and accuracy of two magnetic resonance venography (MRV) methods in a consecutive series of patients with suspected UEDVT.	<ul style="list-style-type: none"> The sensitivity and specificity of time of flight MRV vs gadolinium 3D-MRV was 71% and 89% vs 50% and 80%, respectively. A high number of patients were unable to undergo MRV in this setting. Contrast-enhanced MRV did not improve diagnostic accuracy. The clinical utility of MRV in the setting of suspected UEDVT seems disappointing. 	2

Suspected Upper Extremity Deep Vein Thrombosis
EVIDENCE TABLE

Reference	Study Type	Patients/ Events	Study Objective (Purpose of Study)	Study Results	Strength of Evidence
33. Finn JP, Zisk JH, Edelman RR, et al. Central venous occlusion: MR angiography. <i>Radiology</i> 1993; 187(1):245-251.	9	30	To evaluate time-of-flight MRA in patients with suspected thoracic venous occlusion by comparing MR studies with contrast venography results.	Correlation was excellent between findings of venous obstruction and occlusion at contrast venography and MRA. MRI provided more comprehensive information than catheter venography on central venous anatomy and blood flow. For evaluation of central veins, MRA is an accurate and graphic technique that may succeed in cases in which other methods may give inadequate findings or may be impossible to perform.	3
34. Vogt FM, Herborn CU, Goyen M. MR venography. <i>Magn Reson Imaging Clin N Am</i> 2005; 13(1):113-129, vi.	12	N/A	To review existing MR techniques used for the assessment of the venous system and summarize the clinical experience. New applications that are likely to increase the use of MRV in clinical practice are also reviewed.	Although MRV has successfully shown its potential to evaluate DVT, central venous pathology, and anatomic variants with high accuracy, technical limitations, high cost, limited availability, and logical constraints still prevent its widespread clinical use.	3
35. Cantwell CP, Cradock A, Bruzzi J, Fitzpatrick P, Eustace S, Murray JG. MR venography with true fast imaging with steady-state precession for suspected lower-limb deep vein thrombosis. <i>J Vasc Interv Radiol</i> 2006; 17(11 Pt 1):1763-1769.	9	24	Prospective study to compare true fast imaging with steady-state precession (FISP) MRV for suspected DVT with contrast agent-enhanced venography.	<ul style="list-style-type: none"> When results were analyzed on a per-patient basis, there was good agreement between contrast venography and MRV (kappa=0.64; 95% CI, 0.33-0.94; $p=.0001$). When the venous system was analyzed on a segmental basis, there was very good agreement between contrast venography and MRV (kappa=0.81; 95% CI, 0.68-0.94; $p=.0001$). The sensitivity and specificity for DVT detection were 100% for the iliac and popliteal segments and 100% and 98%, 68% and 94%, and 87% and 98%, respectively, for the femoral, below-knee, and all veins. MRV with axial true FISP allows noninvasive rapid diagnosis of acute DVT in the iliac, femoral, popliteal, and calf muscle veins. MRV is much less reliable in the tibial or peroneal veins. It may demonstrate a nonvenous cause of a patient's symptoms. 	2
36. Miyazaki M, Sugiura S, Tateishi F, Wada H, Kassai Y, Abe H. Non-contrast-enhanced MR angiography using 3D ECG-synchronized half-Fourier fast spin echo. <i>J Magn Reson Imaging</i> 2000; 12(5):776-783.	10	16	To report findings for patients examined with a non-contrast-enhanced MRA using 3D ECG-synchronized half-Fourier fast spin echo.	In the thoracic and abdominal regions, good-quality 3D MRA images were obtained. Furthermore, the weighted subtraction of two images in different phases provides contrast enhancement between arteries and veins.	3

* See Last Page for Key

**Suspected Upper Extremity Deep Vein Thrombosis
EVIDENCE TABLE**

Reference	Study Type	Patients/ Events	Study Objective (Purpose of Study)	Study Results	Strength of Evidence
37. Denson K, Morgan D, Cunningham R, et al. Incidence of venohhhhus thromboembolism in patients with traumatic brain injury. <i>Am J Surg</i> 2007; 193(3):380-383; discussion 383-384.	13	5,787	To review the incidence of venous thromboembolism in critically injured patients with an isolated traumatic brain injury using a standardized venous duplex color-flow Doppler imaging program and to compare it with the overall and high-risk trauma populations.	The incidence of venous thromboembolism in isolated traumatic brain injury is greatest in patients with intraparenchymal hemorrhage. Early venous thromboembolism prophylaxis is warranted in traumatic brain injury patients.	2
38. Tanju S, Sancak T, Dusunceli E, Yagmurlu B, Erden I, Sanlidilek U. Direct contrast-enhanced 3D MR venography evaluation of upper extremity deep venous system. <i>Diagn Interv Radiol</i> 2006; 12(2):74-79.	9	19	To compare MRV and conventional angiograms to determine the diagnostic value of direct contrast-enhanced 3D MRV in mapping the UEDVT and to plan potential interventional procedures.	Results of MRV and conventional angiography were consistent with each other (100% sensitivity and 100% specificity). Direct contrast-enhanced 3D MRV is a well-tolerated sensitive technique in explaining the cause of the malfunctioning arterio-venous fistulas and in pre-surgical planning before placing new catheters or creating fistulas.	3
39. Kim CY, Mirza RA, Bryant JA, et al. Central Veins of the Chest: Evaluation with Time-resolved MR Angiography. <i>Radiology</i> 2008; 247:Epub Mar 18, 2008.	10	27	To retrospectively assess the diagnostic performance of time-resolved MRA in the detection of stenoses and occlusions in the central veins of the chest, with angiographic and surgical findings and consensus readings serving as the reference standard.	Use of time-resolved angiographic data sets as a stand-alone technique had high sensitivity (0.95) but only moderate specificity (0.56) in the detection of venous stenoses or occlusions. Time-resolved angiographic images are a useful adjunct to high-spatial-resolution images in the evaluation of central venous stenoses and occlusions.	2
40. Ruehm S, Kroeger K, Bosk S, Massing S, Mteiescu S, Debatin J, et al. Thromboembolic disease: Assessment with whole body MR venography. <i>Academic Radiology</i> 2005; 12(5, Supplement 1):S63.	9	30	To assess the diagnostic accuracy of a comprehensive MR protocol comprising MRA of the pulmonary arteries and whole body MRV for the detection of thromboembolic disease.	For the detection of venous thrombosis, MRV showed an overall sensitivity and specificity of 95% and 93%, respectively.	2
41. Panzironi G, Rainaldi R, Ricci F, Casale A, De Vargas Macchiucca M. Gray-scale and color Doppler findings in bilateral internal jugular vein thrombosis caused by anaplastic carcinoma of the thyroid. <i>J Clin Ultrasound</i> 2003; 31(2):111-115.	14	1	To report the sonographic findings in a case of bilateral internal jugular vein thrombosis with mild symptoms.	There was evidence of direct infiltration of anaplastic thyroid carcinoma into the left internal jugular vein. Sonographic and CT examinations also demonstrated multiple dilated collateral veins.	4
42. Stam J. Thrombosis of the cerebral veins and sinuses. <i>N Engl J Med</i> 2005; 352(17):1791-1798.	12	N/A	To review pathogenesis of sinus thrombosis, risk factors, clinical and radiologic diagnosis and current evidence and controversies about the best treatment.	Sinus thrombosis remains a diagnostic challenge and a potentially disabling or lethal disease, but improved diagnosis and treatment now result in an excellent outcome for most patients.	4

Suspected Upper Extremity Deep Vein Thrombosis
EVIDENCE TABLE

Reference	Study Type	Patients/ Events	Study Objective (Purpose of Study)	Study Results	Strength of Evidence
43. Kim HC, Chung JW, Yoon CJ, et al. Collateral pathways in thoracic central venous obstruction: three-dimensional display using direct spiral computed tomography venography. <i>J Comput Assist Tomogr</i> 2004; 28(1):24-33.	10	N/A	To illustrate the spectrum of venous collateral pathways caused by the thoracic central venous obstruction with direct spiral CT venography and 3D reconstruction images.	Venous structures that can be demonstrated with CT venography include the jugular veins; the subclavian and brachiocephalic (innominate) veins; the internal and lateral thoracic veins; the superior and inferior venae cavae; the pericardiophrenic veins; the azygos, hemiazygos, and accessory hemiazygos veins; and the intercostal veins	4
44. Sabharwal R, Boshell D, Vladica P. Multidetector spiral CT venography in the diagnosis of upper extremity deep venous thrombosis. <i>Australas Radiol</i> 2007; 51 Suppl:B253-256.	14	1	To present a case which demonstrates the limitations in diagnostic accuracy of US in the diagnosis of UEDVT.	Multidetector spiral CT continues to evolve as a state of the art imaging modality and this case demonstrates the role and value of multidetector spiral CT venography in the diagnosis of UEDVT.	4
45. Kim H, Chung JW, Park JH, et al. Role of CT venography in the diagnosis and treatment of benign thoracic central venous obstruction. <i>Korean J Radiol</i> 2003; 4(3):146-152.	9	18	Prospective study. CT venography and digital subtraction venography were compared to evaluate the role of CT venography in the diagnosis and treatment of benign thoracic central venous obstruction.	The findings of CT venography correlated closely with those of digital subtraction venography. CT venography accurately depicted the degree and extent of benign venous obstruction.	3
46. New CT Protocol Yields Improved Venous Images. <i>RSNA News</i> 2008; 18(1):6-7.	12	N/A	To review a new CT protocol in the diagnosis of venous diseases.	Two researchers, through a trial and error method, have found that changing the contrast bolusing and acquisition timing in multidetector CT of the chest satisfies the demand of thoracic and vascular surgeons for reliable venous images and may also increase reliability in the diagnosis of venous diseases.	3
47. American College of Radiology. <i>Manual on Contrast Media</i> . Available at: http://www.acr.org/SecondaryMainMenu/Categories/quality_safety/contrast_manual.aspx .	15	N/A	Guidance document on contrast media to assist radiologists in recognizing and managing risks associated with the use of contrast media.	N/A	3

Evidence Table Key

Study Type Key

Numbers 1-7 are for studies of therapies while numbers 8-15 are used to describe studies of diagnostics.

1. Randomized Controlled Trial — Treatment
2. Controlled Trial
3. Observation Study
 - a. Cohort
 - b. Cross-sectional
 - c. Case-control
4. Clinical Series
5. Case reviews
6. Anecdotes
7. Reviews

8. Randomized Controlled Trial — Diagnostic
9. Comparative Assessment
10. Clinical Assessment
11. Quantitative Review
12. Qualitative Review
13. Descriptive Study
14. Case Report
15. Other (Described in text)

Strength of Evidence Key

- Category 1 - The conclusions of the study are valid and strongly supported by study design, analysis and results.
- Category 2 - The conclusions of the study are likely valid, but study design does not permit certainty.
- Category 3 - The conclusions of the study may be valid but the evidence supporting the conclusions is inconclusive or equivocal.
- Category 4 - The conclusions of the study may not be valid because the evidence may not be reliable given the study design or analysis.